

University of Groningen

AcmD, a homolog of the major autolysin AcmA of *Lactococcus lactis*, binds to the cell wall and contributes to cell separation and autolysis

Visweswaran, Ganesh Ram R; Steen, Anton; Leenhouts, Kees; Szeliga, Monika; Ruban, Beata; Hesselings-Meinders, Anne; Dijkstra, Bauke W; Kuipers, Oscar P; Kok, Jan; Buist, Girbe

Published in:
 PLoS ONE

DOI:
 [10.1371/journal.pone.0072167](https://doi.org/10.1371/journal.pone.0072167)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Visweswaran, G. R. R., Steen, A., Leenhouts, K., Szeliga, M., Ruban, B., Hesselings-Meinders, A., Dijkstra, B. W., Kuipers, O. P., Kok, J., & Buist, G. (2013). AcmD, a homolog of the major autolysin AcmA of *Lactococcus lactis*, binds to the cell wall and contributes to cell separation and autolysis. *PLoS ONE*, 8(8), [e72167]. <https://doi.org/10.1371/journal.pone.0072167>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Supporting Information




Figure S1.

Comparison of 2D-gel images of supernatant fractions of *L. lactis* IL1403acmA::ISS1 and *L. lactis* IL1403acmA::ISS1acmD::myc. The amount of protein loaded in both cases was the equivalent of supernatant fraction of 100 ml of a GM17 culture with an optical density at 600 nm of 1.0. The position of the spots of the AcmD and AcmD::myc proteins, identified by Mass-spectroscopic analysis, and their molecular weights and pIs are indicated. Proteins that were more abundant in the supernatant fraction of IL1403acmA::ISS1 (blue) or IL1403acmA::ISS1acmD::myc (red), and those unique in the supernatant of IL1403acmA::ISS1 (yellow) or IL1403acmA::ISS1acmD::myc (green) are indicated. Sizes of the pre-stained molecular mass marker (kDa) are indicated in the middle.




Figure S2.

Deletion of *acmD* does not affect cell lysis during growth. Release of intracellular X-prolyl dipeptidyl aminopeptidase (PepX) from *L. lactis* IL1403 (◆), IL1403*acmA::ISS1acmD::myc* (▲) and IL1403*acmA::ISS1* (■). Samples were taken at the indicated time points from the bacterial cultures incubated in GM17 broth. Upon removal of the cells by centrifugation the PepX-activity (in arbitrary units) released into the medium due to autolysis was determined using a chromogenic substrate, as described in the Materials and Methods section




Figure S3.

Expression of $\text{LysM}_{\text{AcmD}}\text{-GFP-His}_{10}$ in *E. coli*. Coomassie brilliant blue-stained SDS- (15%) PAA-gel (left) and *in-gel* GFP- fluorescence (right) showing the expression of $\text{LysM}_{\text{AcmD}}\text{-GFP-His}_{10}$ on SDS- PAGE with a 15% PAA gel. *E. coli* MC1061 bearing the pBADcLIC- $\text{LysM}_{\text{AcmD}}$ was grown at 37° C until OD_{600} of 0.8 and induced with 0.2% arabinose for 2 h (see Materials and Methods section). The cell extracts of control and test samples were loaded on PAA gel for the identification of specific protein band. For the latter figure, the PAA gel is exposed to UV-light prior to coomassie staining for imaging the fluorescent bands. Prestained protein marker lane 1, cell extracts of empty vector control strain, un-induced control and 0.2%-arabinose induced test samples in lanes 2, 3 and 4, respectively. Arrows indicate $\text{LysM}_{\text{AcmD}}\text{-GFP-His}_{10}$ protein/activity bands.




Figure S4.

Negative and autofluorescence controls Phase-contrast and fluorescence microscopy of *L. lactis* NZ9000 cells incubated at pH 4.0 with HIC-purified GFP (a) and without addition of any recombinant protein (b). Original magnification: 1250-fold in all frames.




Figure S5.

Binding of LysM_{AcmD}-GFP-His₁₀ to *L. lactis* NZ9000 cells at pH 6.0 and 8.0. Phase-contrast and fluorescence microscopy of *L. lactis* NZ9000 cells incubated with LysM_{AcmD}-GFP-His₁₀ at pH 6.0 (a) and 8.0 (b). Original magnification: 1250-fold in all frames.